

LAPLACIANS ON QUOTIENTS OF CAUCHY-RIEMANN COMPLEXES AND SZEGÖ MAP FOR L^2 -HARMONIC FORMS

BENT ØRSTED AND GENKAI ZHANG

Dedicated to Professor Gerrit van Dijk on the occasion of his 65th birthday

ABSTRACT. We compute the spectra of the Tanaka type Laplacians $\square = \bar{\partial}_{\mathcal{Q}}^* \bar{\partial}_{\mathcal{Q}} + \bar{\partial}_{\mathcal{Q}} \bar{\partial}_{\mathcal{Q}}^*$ and $\triangle = \partial_{\mathcal{Q}}^* \partial_{\mathcal{Q}} + \partial_{\mathcal{Q}} \partial_{\mathcal{Q}}^*$ on the Rumin complex $\mathcal{Q}$, a quotient of the tangential Cauchy-Riemann complex on the unit sphere S^{2n-1} in $\mathbb{C}^n$. We prove that Szegő map is a unitary operator from a subspace of $(p, q-1)$ -forms on the sphere defined by the operators $\triangle$ and the normal vector field onto the space of L^2 -harmonic (p, q) -forms on the unit ball. Our results generalize earlier result of Folland.

1. INTRODUCTION

The Bergman space of holomorphic functions on the unit ball B of $\mathbb{C}^n$ plays an important role both in representation theory of the group $SU(n, 1)$ and complex analysis. The holomorphic functions in the space can be identified with square integrable harmonic $(n, 0)$ -forms on B equipped with the Bergman metric. One may naturally consider the (p, q) -forms on the unit ball. It has been proved by Donnelly and Fferman [4] that the space of L^2 harmonic (p, q) forms on a strongly pseudo-convex domain is non-empty precisely when $p + q = n$; the case of unit ball was proved earlier in representation theory of semisimple Lie groups and is of much importance for the general strongly pseudo-convex domains. The study of L^2 harmonic (p, q) -forms on the unit ball is already quite involved, partly because the existence theorem does not give explicit construction. In the case of the real ball in $\mathbb{R}^{2n}$ with the hyperbolic metric the L^2 -harmonic n -forms (with respect to the corresponding Laplacian) are Euclidean harmonic forms. One can thus consider the boundary value map of the harmonic forms on the unit sphere, its dual map gives then the Szegő map from $(n-1)$ -form to harmonic n -forms on the real ball. The Szegő map is special case of the general Poisson transform mapping the forms on the sphere to eigenforms on the ball of invariant differential operators. In [6] Gaillard computed explicitly the Poisson transform from $(p-1)$ -forms on the boundary to harmonic (not necessarily L^2) p -forms on the ball. Branson [1], Chen [3], and Lott [9] have found formulas expressing the L^2 -norm of the harmonic forms in terms of their boundary values and in terms of the Laplacian operators on the boundary. These formulas turn out to be quite important both in

Key words and phrases. Differential forms, Cauchy-Riemann complexes, Rumin complexes, Tanaka-Rumin type Laplacians, L^2 -harmonic forms, Szegő map, semisimple Lie groups, unitary representations.

Research by Genkai Zhang supported by the Swedish Science Council (VR) .

conformal geometry, K -theory and in representation theory, see loc. cit. Part of our purpose is to find a corresponding formula for the unit complex ball.

In a remarkable paper [8] Julg and Kasparov give a geometric construction of L^2 -harmonic (p, q) -forms, $p + q = n$, on the unit ball and of the Szegő map from forms on the sphere into harmonic forms. On the sphere S there is the Cauchy-Riemann complex defined as space of $(p, q - 1)$ -forms modulo the ideal generated by the contact form $\bar{\tau} = \sum_{j=1}^n \bar{z}_j dz_j$. However it is proved in [8] that, roughly speaking, the boundary forms of L^2 harmonic (p, q) -forms are in the so-called Rumin complex $\mathcal{Q}$, which is the quotient of Cauchy-Riemann complex modulo the forms τ and $d\tau = \sum_{j=1}^n d\bar{z}_j \wedge dz_j$. In the middle of the Rumin complex is the space of $n - 1$ -forms and there defined a second-order Rumin operator in place of the the first order boundary C-R operator $\bar{\partial}_b$, the boundary value of harmonic forms are in certain Sobolev-type space defined by the Rumin Laplacian which is of 4-order. In the present paper we will use their results to prove a formula expressing the L^2 -norm of the harmonic forms in terms certain Sobolev norm of their boundary values defined in terms of the Tanaka-type Laplacians $\square_{\mathcal{Q}} = \bar{\partial}_{\mathcal{Q}}^* \bar{\partial}_{\mathcal{Q}} + \bar{\partial}_{\mathcal{Q}} \bar{\partial}_{\mathcal{Q}}^*$ and $\triangle_{\mathcal{Q}} = \partial_{\mathcal{Q}}^* \partial_{\mathcal{Q}} + \partial_{\mathcal{Q}} \partial_{\mathcal{Q}}^*$ on the complex $\mathcal{Q}$ on the unit sphere. For that purpose we compute the spectrum of the Laplacians on the Rumin complex of (p, q) -forms for general (p, q) . For $(0, q)$ -forms the operator $\square_{\mathcal{Q}}$ coincides with the usual C-R Laplacian $\square_b$ and its spectrum has been found earlier by Folland [5]. So our results generalize that of Folland.

The L^2 -space of differential forms in the Rumin complex on the sphere is a natural model for the induced representations of the group $SU(1, n)$ from certain finite-dimensional representations of the maximal parabolic subgroup. There arise many other geometrically interesting differential operators and unitary representations such as complementary series, see e.g.[2]. We hope that our results here will shed light on studying those subjects.

The paper is organized as follows. In Section 2 we present the context of our problems and recall the known results. In Section 3 we compute the the eigenvalues of the Laplacians $\square_b$ and $\triangle_b$ on certain subspaces of $(p, q - 1)$ forms. The unitary structure of the space of L^2 harmonic (p, q) -forms is given in Section 4.

We would like to thank the organizers of the conference “Harmonic Analysis and Homogeneous Spaces”, Lorentz Center, Leiden University, August 23-27, 2004, for the invitation. Thanks are also due to Thomas Branson for some helpful discussions and to Miraslav Engliš for reading an earlier version of this paper.

2. CAUCHY-RIEMANN COMPLEXES AND THEIR QUOTIENTS

For some notational convenience we will consider $(p, q - 1)$ -forms on the unit sphere S . We assume throughout the text that $n > 1$, $1 \leq p, q \leq n$.

We recall first briefly some well-known facts about Cauchy-Riemann complex of differential forms on S .

Let $\Omega^{p, q-1}$ be the space of smooth differential $(p, q - 1)$ -forms on some neighborhood of the unit sphere S . Recall by definition that (p, q) -form is a section of the vector bundle $\wedge^q T^{(0,1)'} \otimes \wedge^p T^{(1,0)'}$ over some neighborhood of S . For simplicity we denote

also $\Omega_z^{p,q-1} = \wedge^{q-1} T_z^{(0,1)'} \otimes \wedge^p T_z^{(1,0)'}$ the fiber space at each z . We fix also the usual convention that the $(1,0)$ -covectors in $T_z^{(1,0)'}$ and the $(0,1)$ -covectors in $T_z^{(0,1)'}$ are anticommuting.

We let

$$\tau = \sum_{j=1}^n \bar{z}_j dz_j, \quad \bar{\tau} = \sum_{j=1}^n z_j d\bar{z}_j,$$

be the $(1,0)$ - and $(0,1)$ -forms defining the contact structure on $S = \partial B$. We let $\Omega^{p,q-1}/(\bar{\tau})$ be the quotient by the ideal $(\bar{\tau})$ generated by $\bar{\tau}$. Since $\bar{\partial}\bar{\tau} = 0$, there is an induced operator, denoted by $\bar{\partial}_b$ on the quotient $\Omega^{p,q-1}/(\bar{\tau})$ and it forms the C-R complex.

For our purpose we need to consider some further quotients. Let $\mathcal{Q}^{p,q-1} = \Omega^{p,q-1}/(\tau, \bar{\tau}, d\tau)$ be the quotient of $\Omega^{p,q-1}$ by the ideal generated by τ , $\bar{\tau}$ and $d\tau$; the ideal is clearly invariant under ∂ and $\bar{\partial}$, the induced operators on the quotient $\mathcal{Q}^{p,q-1}$ will be denoted by $\partial_{\mathcal{Q}}$ and $\bar{\partial}_{\mathcal{Q}}$. The spaces $\sum_{p+q-1=j} \mathcal{Q}^{p,q-1}$ for $j \leq n-1$ and the induced operator of $d = \bar{\partial} + \partial$ can be prolonged to a complex, also called Rumin complex, so that the cohomology so obtained is the usual de Rham cohomology, see [8] and references therein.

Consider the unitary group $U(n)$ acting natural on tangent space $\mathbb{C}^n = \sum_{j=1}^n \mathbb{C} \frac{\partial}{\partial z_j}$ of B by definition, and its dual action on the differential forms $\sum_{j=1}^n \mathbb{C} dz_j$ of degree 1 with constant coefficients. We fix the unique $U(n)$ -invariant point-wise inner product $\langle \cdot, \cdot \rangle$ on the differential forms at each point normalized so that $d\bar{z}_{j_1} \wedge \cdots \wedge d\bar{z}_{j_q} \wedge dz_{i_1} \wedge \cdots \wedge dz_{i_p}$, $1 \leq i_1 < \cdots < i_p \leq n$, $1 \leq j_1 < \cdots < j_q$ form an orthonormal basis of different (p, q) forms at each $z \in S$. We define the $U(n)$ -invariant inner product on $\Omega^{p,q-1}$ by

$$(\alpha, \beta) = \int_S \langle \alpha(z), \beta(z) \rangle d\sigma(z)$$

where $d\sigma(z)$ is the normalized area measure on S .

We need further some important operators on differential forms. For $\alpha \in \Omega^{p,q-1}$ we denote L_α the corresponding multiplication operators by α , $L_\alpha \beta = \alpha \wedge \beta$. For a holomorphic tangent or anti-holomorphic tangent vector X in $\mathbb{C}^n$ we let i_X be the standard evaluation operator see e.g. [7, Chapter 0, Section 7]. In particular the operators $\partial, L_{dz_j}, i_{\partial_j}$ anti-commutes with the operators $\bar{\partial}, L_{d\bar{z}_j}, i_{\bar{\partial}_j}$. Let

$$T = \sum_{j=1}^n z_j \partial_j, \quad \bar{T} = \sum_{j=1}^n \bar{z}_j \bar{\partial}_j,$$

and

$$E = \sum_{j=1}^n \partial_j \otimes \bar{\partial}_j$$

and i_E the corresponding evaluation operator

$$i_E = \sum_{j=1}^n i_{\partial_j} i_{\bar{\partial}_j}.$$

Viewed as operators on $\Omega^{p,q-1}$ with the point-wise inner product $\langle \alpha(z), \beta(z) \rangle$ for each point z their adjoint operators are

$$(2.1) \quad (L_{d\tau})^* = i_E, \quad (L_\tau)^* = i_T, \quad (L_{\bar{\tau}})^* = i_{\bar{T}}.$$

This is then also true with respect to the inner product (α, β) .

The space $\mathcal{Q}^{p,q-1} = \Omega^{p,q-1}/(\tau, \bar{\tau}, d\tau)$ can be realized as the subbundle $\text{Ker } i_{\bar{T}} \cap \text{Ker } i_T \cap \text{Ker } i_E$ on $\Omega^{p,q-1}$ and we will fix this realization throughout our paper. The induced C-R operators ∂_Q and $\bar{\partial}_Q$ will be, under this identification, the differentiations ∂ and $\bar{\partial}$, followed by the projection onto the sub-bundle. We put

$$(2.2) \quad \zeta_j := \bar{\partial}_b \bar{z}_j = \bar{\partial}_Q \bar{z}_j = d\bar{z}_j - \bar{z}_j \bar{\tau}, \quad \xi_j = \partial_Q z_j = dz_j - z_j \tau;$$

see e.g. Folland [5].

The next lemma is elementary.

Lemma 2.1. The differential form $d\tau - \bar{\tau} \wedge \tau$ is T and $\bar{T}$ -primitive, namely it is annihilated by i_T and $i_{\bar{T}}$. Moreover the forms ζ_j and ξ_j can also be obtained by

$$\zeta_j = -i_{\partial_j}(d\tau - \bar{\tau} \wedge \tau), \quad \xi_j = i_{\bar{\partial}_j}(d\tau - \bar{\tau} \wedge \tau).$$

We need also some Hodge type commutation relations; the first equality below is well known (see e.g. [7, Chapter 0, Section 7]), the other can be proved by routine computation.

Lemma 2.2. Assume $p + q \leq n$. The following commutation relations hold, viewed as operators acting on the space $\Omega^{p,q-1}$,

$$\begin{aligned} [i_E, L_{d\tau}] &= p + q \\ [i_T, L_{d\tau}] &= L_{\bar{\tau}}, \quad [i_{\bar{T}}, L_{d\tau}] = L_\tau \end{aligned}$$

and

$$i_T L_\tau + L_\tau i_T = |z|^2, \quad i_{\bar{T}} L_{\bar{\tau}} + L_{\bar{\tau}} i_{\bar{T}} = |z|^2.$$

By taking adjoint we get then

$$(2.3) \quad [i_E, L_\tau] = -i_T, \quad [i_E, L_{\bar{\tau}}] = i_{\bar{T}}.$$

It follows further that

$$(2.4) \quad [i_T, L_{d\tau - \bar{\tau} \wedge \tau}] = 0,$$

and

$$(2.5) \quad [i_E - i_{\bar{T}} i_T, L_{d\tau - \bar{\tau} \wedge \tau}] = (n-1) - \sum_{j=1}^n L_{\zeta_j} i_{\partial_j} - \sum_{j=1}^n L_{\xi_j} i_{\bar{\partial}_j}.$$

The following decomposition, which will not be used in this paper, follows easily from our previous commutation relation, and it might be of some independent interests; it can be proved easily and we omit the proof.

Corollary 2.3. Realizing the quotient $\mathcal{Q}^{p,q-1}$ as the space $\text{Ker } i_T \cap \text{Ker } i_{\bar{T}} \cap \text{Ker } i_E$ on $\Omega^{p,q-1}$, we have

$$\Omega^{p,q-1} = \mathcal{Q}^{p,q-1} \oplus (L_\tau \Omega^{p-1,q-1} + L_{\bar{\tau}} \Omega^{p-1,q-1}) \oplus L_{d\tau - \bar{\tau} \wedge \tau} \Omega^{p-1,q-2}$$

To state the next result we fix the space $\mathfrak{h} = \mathbb{C}H_1 + \cdots + \mathbb{C}H_n$, of diagonal matrices as a Cartan subalgebra of the Lie algebra $\mathfrak{u}(n)^\mathbb{C} = \mathfrak{gl}(n, \mathbb{C})$, with $H_j = \text{diag}(0, \dots, 0, 1, 0, \dots, 0)$, $j = 1, \dots, n$. Let $\{\varepsilon_j\}$ be the basis of $\mathfrak{h}'$ dual to $\{H_j\}$, and we fix an ordering of roots of $\mathfrak{h}$ in $\mathfrak{gl}(n, \mathbb{C})$ so that $\varepsilon_1 > \cdots > \varepsilon_n$.

Proposition 2.4. Assume $p + q \leq n$. As a representation space of $K = U(n)$ the L^2 -space of $(p, q-1)$ -forms in the quotient $\mathcal{Q}^{p, q-1} = \Omega^{p, q}/(\tau, \bar{\tau}, d\tau)$ is decomposed with multiplicity free as follows

$$L^2(\mathcal{Q}^{p, q-1}) = \sum_{m, l \geq 0}^{\oplus} (\Phi_{q-1, p}^{m, l} \oplus \Phi_{q-1, p+1}^{m, l} \oplus \Phi_{q, p}^{m, l} \oplus \Phi_{q, p+1}^{m, l}),$$

where $\Phi_{r, s}^{m, l}$ are of highest weight

$$(m, 0, \dots, 0, -l) + (\overbrace{1, \dots, 1}^r, 0, \dots, 0, \overbrace{-1, \dots, -1}^s)$$

with r many 1's and s many -1's, for $(r, s) = (q-1, p), (q-1, p+1), (q, p), (q, p+1)$; the last summand will not appear when $p + q = n$.

Proof. This follows from the Frobenius reciprocity and is stated in [8] (without proof). We give a brief proof here. View the space S as a homogeneous space $S = U(n)/U(n-1)$ where $U(n-1)$ is the subgroup keeping the base point $z_0 = (1, 0, \dots, 0) \in S$ fixed. The bundle $\mathcal{Q}^{p, q-1}$ is a $U(n)$ -homogeneous bundle, and the fiber space at the base point is spanned by the vectors $d\bar{z}_{i_1} \wedge \cdots \wedge d\bar{z}_{i_{q-2}} \wedge dz_{j_1} \wedge \cdots \wedge dz_{j_{p-1}}$, $2 \leq i_1 < \cdots < i_{q-1} \leq n$, $2 \leq j_1 < \cdots < j_{p-1} \leq n$ modulo the ideal generated by the form $d\bar{z}_2 \wedge dz_2 + \cdots + d\bar{z}_n \wedge dz_n$ since $\tau(z_0) = dz_1$ and $\bar{\tau}(z_0) = d\bar{z}_1$, and it forms an irreducible representation ω_0 of $U(n-1)$, by e.g. the Lefschetz decomposition [7] and by the tensor product decomposition of representations of $U(n-1)$ [11]. The multiplicity of a representation μ of $U(n)$ appearing in $L^2(\mathcal{Q}^{p, q-1})$ is the same as the multiplicity of ω_0 in μ . The rest follows then by the known results on branching under $U(n-1)$ of representations of $U(n)$, see e.g. [11]. $\square$

3. SPECTRUM OF THE LAPLACIANS $\square_{\mathcal{Q}} = \bar{\partial}_{\mathcal{Q}}^* \bar{\partial}_{\mathcal{Q}} + \bar{\partial}_{\mathcal{Q}} \bar{\partial}_{\mathcal{Q}}^*$ AND $\triangle_{\mathcal{Q}} = \partial_{\mathcal{Q}}^* \partial_{\mathcal{Q}} + \partial_{\mathcal{Q}} \partial_{\mathcal{Q}}^*$

In this section we compute the spectrum of the Tanaka-type Laplacians $\square_{\mathcal{Q}}$ and $\triangle_{\mathcal{Q}}$ in the space $L^2(\mathcal{Q}^{p, q-1})$, by computing their eigenvalues on each irreducible subspace. To simplify the notation we will suppress the upper indices (m, l) .

Lemma 3.1. The highest weight vectors $\alpha, \beta, \gamma, \delta$ in the spaces $\Phi_{q-1, p}^{m, l}$, $\Phi_{q-1, p+1}^{m, l}$, $\Phi_{q, p}^{m, l}$ and $\Phi_{q, p+1}^{m, l}$ are given as follows:

$$\alpha_{q-1, p} = \bar{z}_1^m z_n^l \alpha'_{q-1, p}$$

with

$$\begin{aligned} \alpha'_{q-1, p} = & \zeta_1 \wedge \cdots \wedge \zeta_{q-1} \wedge \xi_{n-p+1} \wedge \cdots \wedge \xi_n \\ & - \frac{1}{n-1} (d\tau - \bar{\tau} \wedge \tau) \wedge i_E(\zeta_1 \wedge \cdots \wedge \zeta_{q-1} \wedge \xi_{n-p+1} \wedge \cdots \wedge \xi_n), \end{aligned}$$

$$\beta_{q-1,p+1} = \bar{z}_1^m z_n^l \zeta_1 \wedge \cdots \wedge \zeta_{q-1} \wedge \left(\sum_{i=n-p}^n (-1)^{i-(n-p)} z_i \xi_{n-p} \wedge \cdots \widehat{\xi_i} \cdots \wedge \xi_n \right),$$

$$\gamma_{q,p} = \bar{z}_1^m z_n^l \left(\sum_{j=1}^q (-1)^{j-1} \bar{z}_j \zeta_1 \wedge \cdots \widehat{\zeta_j} \cdots \wedge \zeta_q \right) \wedge \xi_{n-p+1} \wedge \cdots \wedge \xi_n$$

and respectively

$$\delta_{q,p+1} = \bar{z}_1^m z_n^l \left(\sum_{j=1}^q (-1)^{j-1} \bar{z}_j \zeta_1 \wedge \cdots \widehat{\zeta_j} \cdots \wedge \zeta_q \right) \wedge \left(\sum_{i=n-p}^n (-1)^{i-(n-p)} z_i \xi_{n-p} \wedge \cdots \widehat{\xi_i} \cdots \wedge \xi_n \right).$$

Proof. It is easy to prove that the given forms are of the respective weights. By the multiplicity free result in Proposition 2.4 we need only to prove that they are in $\mathcal{Q}$. We consider first the forms β, γ, δ . It follows from [5, Theorem 5] that

$$(3.1) \quad \omega_1(q) := \sum_{j=1}^q (-1)^{j-1} \bar{z}_j \zeta_1 \wedge \cdots \widehat{\zeta_j} \cdots \wedge \zeta_q = \sum_{j=1}^q (-1)^{j-1} \bar{z}_j d\bar{z}_1 \wedge \cdots \widehat{d\bar{z}_j} \cdots \wedge d\bar{z}_q,$$

and (by rewriting the formula there)

$$(3.2) \quad \zeta_1 \wedge \cdots \wedge \zeta_{q-1} = d\bar{z}_1 \wedge \cdots \wedge d\bar{z}_{q-1} - \bar{\tau} \wedge \omega_1(q-1)$$

and they are highest weight vectors in the C-R complex $\mathcal{Q}^{0,q}$ of $(0, q)$ -forms. Similarly we can prove that

$$(3.3) \quad \begin{aligned} \omega_2(p) &:= \sum_{i=n-p}^n (-1)^{i-(n-p)} z_i \xi_{n-p} \wedge \cdots \wedge \widehat{\xi_i} \cdots \wedge \xi_n \\ &= \sum_{i=n-p}^n (-1)^{i-(n-p)} z_i dz_{n-p} \wedge \cdots \widehat{dz_i} \cdots \wedge dz_n \end{aligned}$$

and

$$(3.4) \quad \xi_{n-p+1} \wedge \cdots \wedge \xi_n = dz_{n-p+1} \wedge \cdots \wedge dz_n - \tau \wedge \omega_2(p-1)$$

are highest weight vectors in the C-R complex $\mathcal{Q}^{p,0}$ of $(p, 0)$ -forms. The forms β, γ and δ are thus all annihilated by i_T and $i_{\bar{T}}$; moreover we have $i_E(\delta) = 0$ by a simple computation. We prove now that $i_E(\gamma) = 0$. We have, by the above formulas,

$$\gamma = \omega_1(q) \wedge dz_{n-p+1} \wedge \cdots \wedge dz_n - (-1)^{q-1} \tau \wedge \omega_1(q) \wedge \omega_2(p-1).$$

The first term is clearly annihilated by i_E , and so is also $\omega_1(q) \wedge \omega_2(p-1)$. Thus by the commutation relation (2.3), we have that up to a factor of ± 1 , $i_E(\gamma)$ is

$$i_{\bar{T}}(\omega_1(q) \wedge \omega_2(p-1)) = 0$$

since each term $\omega_1(q)$ and $\omega_2(p-1)$ is annihilated by $i_{\bar{T}}$. The proof of $i_E(\beta) = 0$ is the same.

Consider finally the form α . We have that $i_T(\alpha) = 0$, $i_{\bar{T}}(\alpha) = 0$, since the forms ζ 's and ξ 's and $d\tau - \bar{\tau} \wedge \tau$ are annihilated by i_T and $i_{\bar{T}}$. We prove now $i_E(\alpha) = 0$. Write

$$\alpha' = \kappa - \frac{1}{n-1}(d\tau - \bar{\tau} \wedge \tau) \wedge i_E(\kappa)$$

with

$$\kappa = \zeta_1 \wedge \cdots \wedge \zeta_{q-1} \wedge \xi_{n-p+1} \wedge \cdots \wedge \xi_n.$$

Now

$$\begin{aligned} i_E(\alpha') &= i_E(\kappa) - \frac{1}{n-1}i_E((d\tau - \bar{\tau} \wedge \tau)i_E(\kappa)) \\ &= i_E(\kappa) - \frac{1}{n-1}(i_E(d\tau - \bar{\tau} \wedge \tau)i_E(\kappa) + (d\tau - \bar{\tau} \wedge \tau) \wedge i_E i_E(\kappa)) \\ &= i_E(\kappa) - \frac{1}{n-1}(n-1)i_E(\kappa) - \frac{1}{n-1}(d\tau - \bar{\tau} \wedge \tau) \wedge i_E i_E(\kappa) \\ &= -\frac{1}{n-1}(d\tau - \bar{\tau} \wedge \tau) \wedge i_E i_E(\kappa) \end{aligned}$$

since $(i_E(d\tau) - i_E(\bar{\tau} \wedge \tau)) = n-1$. We claim that

$$(3.5) \quad i_E i_E(\kappa) = 0.$$

Indeed, the form κ is

$$\begin{aligned} \kappa &= d\bar{z}_1 \wedge \cdots \wedge d\bar{z}_{q-1} \wedge dz_{n-p+1} \wedge \cdots \wedge dz_n - \bar{\tau} \wedge \omega_1 \wedge dz_{n-p+1} \wedge \cdots \wedge dz_n \\ &\quad - (-1)^{q-1} \tau \wedge d\bar{z}_1 \wedge \cdots \wedge d\bar{z}_{q-1} \wedge \omega_2 + (-1)^{q-1} \bar{\tau} \wedge \tau \wedge \omega_1 \wedge \omega_2 \end{aligned}$$

with $\omega_1 = \omega_1(q-1)$ and $\omega_2 = \omega_2(p-1)$ defined in (3.3)-(3.4) (q and p replaced by $q-1$ and $p-1$ respectively). From this we have

$$\begin{aligned} i_E(\kappa) &= i_E(\bar{\tau} \wedge \omega_1 \wedge dz_{n-p+1} \wedge \cdots \wedge dz_n) - (-1)^{q-1} i_E(\tau \wedge d\bar{z}_1 \wedge \cdots \wedge d\bar{z}_{q-1} \wedge \omega_2) \\ &\quad + (-1)^{q-1} i_E(\bar{\tau} \wedge \tau \wedge \omega_1 \wedge \omega_2). \end{aligned}$$

Using the commutation relation (2.3) and using the fact that all the forms $d\bar{z}_1 \wedge \cdots \wedge d\bar{z}_{q-1} \wedge \omega_2$, $\omega_1 \wedge dz_{n-p+1} \wedge \cdots \wedge dz_n$ and ω_1 and ω_2 are forms in $\mathcal{Q}$ we get

$$(3.6) \quad i_E(\kappa) = (-1)^{q-1} \omega_1 \wedge \omega_2,$$

and (3.5) follows, again by the formulas (3.1) and (3.3). □

The following lemma gives the L^2 -norms of the highest weight vectors α , β , γ , and δ . For computational reasons we assume $p+q < n$ for certain cases. If $p+q = n$ one can find the norm by the same computation which we omit here.

Lemma 3.2. The following formulas hold

$$\begin{aligned} \|\alpha_{q-1,p}\|^2 &= \frac{m!l!}{(n)_{m+l+2}} \\ &\quad \times \left((l+n-q+1)(m+n-p) - \frac{1}{n-1}(l+p)(m+q-1) \right), \quad (p+q < n) \\ \|\beta_{q-1,p+1}\|^2 &= \frac{m!l!}{(n)_{m+l+2}} (l+p+1)(l+n-q), \end{aligned}$$

$$\begin{aligned}\|\gamma_{q,p}\|^2 &= \frac{m!l!}{(n)_{m+l+2}}(m+q)(m+n-p+1), \\ \|\delta_{q,p+1}\|^2 &= \frac{m!l!}{(n)_{m+l+2}}(l+p+1)(m+q). \quad (p+q < n)\end{aligned}$$

Proof. We consider first the forms β , γ and δ . The pointwise inner products $\langle\beta, \beta\rangle$, $\langle\gamma, \gamma\rangle$, and $\langle\delta, \delta\rangle$ can be computed using the equalities (3.1)-(3.4) and the orthogonality of the forms $d\bar{z}_{j_1} \wedge \cdots \wedge d\bar{z}_{j_{q-1}} \wedge dz_{i_1} \wedge \cdots \wedge dz_{i_p}$ (see also [5], though a different normalization was used), we have

$$\begin{aligned}\langle\beta, \beta\rangle &= |\bar{z}_1^m z_n^l|^2 \left(\sum_{i=n-p}^n |z_i|^2 \right) \left(\sum_{j=q}^n |z_j|^2 \right), \\ \langle\gamma, \gamma\rangle &= |\bar{z}_1^m z_n^l|^2 \left(\sum_{i=1}^{n-p} |z_i|^2 \right) \left(\sum_{j=1}^q |z_j|^2 \right), \\ \langle\delta, \delta\rangle &= |\bar{z}_1^m z_n^l|^2 \left(\sum_{j=1}^q |z_j|^2 \right) \left(\sum_{i=n-p}^n |z_i|^2 \right).\end{aligned}$$

Their L^2 -norms can be computed then by integration over S , and we omit the details.

We compute now the norm of α . Adapting the notation in the proof of Lemma 3.1, we have

$$\begin{aligned}\langle\alpha', \alpha'\rangle &= \langle\kappa, \kappa\rangle - \frac{1}{n-1} \langle\kappa, (d\tau - \bar{\tau} \wedge \tau) \wedge i_E \kappa\rangle - \frac{1}{n-1} \langle(d\tau - \bar{\tau} \wedge \tau) \wedge i_E \kappa, \kappa\rangle \\ &\quad + \frac{1}{(n-1)^2} \langle(d\tau - \bar{\tau} \wedge \tau) \wedge i_E \kappa, (d\tau - \bar{\tau} \wedge \tau) \wedge i_E \kappa\rangle.\end{aligned}$$

The inner product in the second term is

$$\langle\kappa, (d\tau - \bar{\tau} \wedge \tau) \wedge i_E \kappa\rangle = \langle(i_E - i_T i_{\bar{T}}) \kappa, i_E \kappa\rangle = \langle i_E \kappa, i_E \kappa\rangle$$

since κ is in $\text{Ker } i_T$. So the third term is also $\langle i_E \kappa, i_E \kappa\rangle$. The inner product in the last term is

$$\begin{aligned}\langle(d\tau - \bar{\tau} \wedge \tau) \wedge i_E \kappa, (d\tau - \bar{\tau} \wedge \tau) \wedge i_E \kappa\rangle &= \langle(i_E - i_T i_{\bar{T}}) ((d\tau - \bar{\tau} \wedge \tau) \wedge i_E \kappa), i_E \kappa\rangle \\ &= \langle i_E ((d\tau - \bar{\tau} \wedge \tau) \wedge i_E \kappa), i_E \kappa\rangle - \langle i_T (i_{\bar{T}} ((d\tau - \bar{\tau} \wedge \tau) \wedge i_E \kappa)), i_E \kappa\rangle.\end{aligned}$$

But by (2.4)

$$i_{\bar{T}}((d\tau - \bar{\tau} \wedge \tau) \wedge i_E \kappa) = (d\tau - \bar{\tau} \wedge \tau) \wedge i_{\bar{T}} i_E \kappa$$

and $i_{\bar{T}} i_E \kappa = i_E i_{\bar{T}} \kappa = 0$. So this inner product is

$$\begin{aligned}\langle i_E ((d\tau - \bar{\tau} \wedge \tau) \wedge i_E \kappa), i_E \kappa\rangle &= (n-1) \langle i_E \kappa, i_E \kappa\rangle + \langle(d\tau - \bar{\tau} \wedge \tau) \wedge (i_E i_E \kappa), i_E \kappa\rangle \\ &= (n-1) \langle i_E \kappa, i_E \kappa\rangle\end{aligned}$$

by (3.5). We have then

$$(3.7) \quad \langle\alpha, \alpha\rangle = |\bar{z}_1^m z_n^l|^2 \langle\kappa, \kappa\rangle - \frac{1}{n-1} |\bar{z}_1^m z_n^l|^2 \langle i_E \kappa, i_E \kappa\rangle.$$

The point-wise norm squared of κ is

$$\langle\kappa, \kappa\rangle = \left(\sum_{i=1}^{n-p} |z_i|^2 \right) \left(\sum_{j=q}^n |z_j|^2 \right).$$

Using (3.6) we find then that

$$\langle i_E(\kappa), i_E(\kappa) \rangle = \langle \omega_1 \wedge \omega_2, \omega_1 \wedge \omega_2 \rangle = \langle \omega_1, \omega_1 \rangle \langle \omega_2, \omega_2 \rangle = \left(\sum_{j=1}^{q-1} |z_j|^2 \right) \left(\sum_{i=n-p+1}^n |z_i|^2 \right).$$

The integrals of the two terms in (3.7) can be computed directly and we prove then our results. $\square$

Theorem 3.3. Assume $p, q \geq 1$ and $p + q \leq n$.

- (1) The spectrum of the Laplacian $\square_{\mathcal{Q}} = \bar{\partial}_{\mathcal{Q}}^* \bar{\partial}_{\mathcal{Q}} + \bar{\partial}_{\mathcal{Q}} \bar{\partial}_{\mathcal{Q}}^*$ on $L^2(\mathcal{Q}^{p,q-1})$ is given as follows: The subspaces $\Phi_{q-1,p}^{m,l}$ ($p+q < n$), $\Phi_{q-1,p+1}^{m,l}$, $\Phi_{q,p}^{m,l}$ ($p+q < n$), and $\Phi_{q,p+1}^{m,l}$ ($p+q < n$) are eigenspaces of $\square_{\mathcal{Q}}$ with eigenvalues

$$\frac{m+q-1}{m+n-p+1} \left((l+n-q+1)(m+n-p) - \frac{1}{n-1} (l+p)(m+q-1) \right),$$

$$(m+q-1)(l+n-q+2), \quad (m+q-1)(l+n-q+1), \quad (m+q)(l+n-q+1)$$

respectively.

- (2) The spectra of the Laplacians $\Delta_{\mathcal{Q}} = \partial_{\mathcal{Q}}^* \partial_{\mathcal{Q}} + \partial_{\mathcal{Q}} \partial_{\mathcal{Q}}^*$ on $L^2(\mathcal{Q}^{p,q-1})$ are given as follows: The subspaces $\Phi_{q-1,p}^{m,l}$ ($p+q < n$), $\Phi_{q-1,p+1}^{m,l}$ ($p+q < n$), $\Phi_{q,p}^{m,l}$, and $\Phi_{q,p+1}^{m,l}$ ($p+q < n$) are eigenspaces of $\Delta_{\mathcal{Q}}$ with eigenvalues

$$\frac{l+p}{l+n-q} \left((l+n-q+1)(m+n-p) - \frac{1}{n-1} (l+p)(m+q-1) \right),$$

$$(l+p)(m+n-p+1), \quad (l+p)(m+l-p+1), \quad (l+p)(m+n-p+1)$$

respectively.

Proof. To simplify the notation we will suppress the subindex $\mathcal{Q}$ and write $\partial = \partial_{\mathcal{Q}}$. The operators ∂ , and $\bar{\partial}$, and their adjoint operators are clearly K -invariant operators. By the multiplicity free result in Proposition 2.4, we have that the operator $\bar{\partial}_{\mathcal{Q}}$ maps highest weight vectors of bi-degree $(p-1, q-1)$ to highest weight vectors of bi-degree $(p, q-1)$, and the corresponding statement is true for the operator $\partial_{\mathcal{Q}}$ and the adjoint operators. Explicitly we have, by direct computations,

$$\bar{\partial}\alpha = 0, \quad \partial\alpha = 0$$

$$\bar{\partial}\beta = 0, \quad \partial\beta_{q-1,p+1} = (l+p+1)\alpha_{q-1,p+1},$$

$$\bar{\partial}\gamma_{q,p} = (m+q)\alpha_{q,p}, \quad \partial\gamma_{q,p} = 0$$

$$\bar{\partial}\delta_{q,p+1}^{m,l} = (m+q)\beta_{q,p+1}^{m,l}, \quad \partial\delta_{q,p+1}^{m,l} = (l+p+1)\gamma_{q,p+1}^{m,l}.$$

By the multiplicity free result we get then

$$\bar{\partial}^* \alpha_{q-1,p} = (m+q-1) \frac{\|\alpha_{q-1,p}\|^2}{\|\gamma_{q-1,p}\|^2} \gamma_{q-1,p}, \quad \partial^* \alpha_{q-1,p} = (l+p) \frac{\|\alpha_{q-1,p}\|^2}{\|\beta_{q-1,p}\|^2} \beta_{q-1,p}$$

$$\bar{\partial}^* \beta_{q-1,p+1} = (m+q-1) \frac{\|\beta_{q-1,p+1}\|^2}{\|\delta_{q-1,p+1}\|^2} \delta_{q-1,p+1}, \quad \partial^* \beta_{q-1,p+1} = 0$$

$$\begin{aligned}\bar{\partial}^* \gamma_{q,p} &= 0, \quad \partial^* \gamma_{q,p} = (l+p) \frac{\|\gamma_{q,p}\|^2}{\|\delta_{q,p}\|^2} \delta_{q,p}, \\ \bar{\partial}^* \delta_{q,p+1} &= 0, \quad \partial^* \delta_{q-1,p+1} = 0.\end{aligned}$$

We compute the eigenvalues of $\square$ on $\alpha_{q-1,p}$,

$$\square \alpha_{q-1,p} = \partial \bar{\partial}^* \alpha_{q-1,p} = (m+q-1) \frac{\|\alpha_{q-1,p}\|^2}{\|\gamma_{q-1,p}\|^2} \partial \gamma_{q-1,p} = (m+q-1)^2 \frac{\|\alpha_{q-1,p}\|^2}{\|\gamma_{q-1,p}\|^2} \alpha_{q-1,p},$$

and our result then follows by Lemma 3.2. The other eigenvalues can be computed similarly. $\square$

4. $SU(1, n)$ -INVARIANT UNITARY STRUCTURE ON THE HARMONIC FORMS VIA THE SZEGÖ MAP AND THE TANAKA LAPLACIANS

In the paper [8] Julg and Kasparov give a geometric construction of the L^2 -harmonic (p, q) -forms on B . We recall some of their results here.

Adapting the notation there we put

$$a = p + l, \quad b = q + m.$$

Let $\omega = \bar{z}_1^l z_n^m d\bar{z}_1 \wedge \cdots \wedge d\bar{z}_q \wedge dz_{n-q+1} \wedge \cdots \wedge dz_n$ be a (p, q) -form on C^n , which is closed and primitive in the Euclidean metric (thus Euclidean harmonic). Let $F(t) = {}_2F_1(a, b; a+b+2, t)$, $t = |z|^2$, be the hypergeometric function. Put

$$\tilde{\omega} = f_0 \omega + f_1 \tau \wedge i_T \omega + f_2 \bar{\tau} \wedge i_{\bar{T}} \omega + f_3 \tau \wedge \bar{\tau} \wedge i_{\bar{T}} i_T \omega$$

where

$$f_0(z) = abF(t) - tF'(t), f_1(z) = (b+1)F'(t), f_2(z) = (a+1)F'(t), f_3(z) = F''(t).$$

Theorem 4.1. [8, Sections 3-4] Let $p, q \geq 1$ and $p+q = n$. The forms $\tilde{\omega}$ are L^2 -harmonic (p, q) -forms and they form an orthogonal basis for the space $\mathcal{H}^{p,q}$. The Szegő map S maps $L^2(\mathcal{Q}^{n-1})$ onto $\mathcal{H}^{p,q}$ and we have

$$S(\gamma_{q,p}^{m,l}) = \frac{1}{aF(1)} \tilde{\omega}.$$

In [8, Theorem 4.9] the unitarity of the Szegő map is studied on the space of all L^2 -harmonic n -forms using the Rumin operator, which is a fourth order differential operator. We prove now a unitarity result using the operator $\partial_{\mathcal{Q}}$ and the vector field $T + \bar{T}$.

Define $W_{p,q-1}^{1,2}$ to be the Hilbert space completion of differential forms in $\Psi_{q,p}^{m,l}$, $m, l \geq 0$ with the norm

$$(\gamma, \gamma)_{W^{1,2}} = ((\partial_{\mathcal{Q}} \partial_{\mathcal{Q}}^* - (T + \bar{T})_*) \gamma, \gamma).$$

Here $(T + \bar{T})_*$ is induced action of vector field $T + \bar{T}$ on differential forms. $W_{p,q-1}^{1,2}$ can be viewed as a Sobolev space of differential forms.

Theorem 4.2. Let $p, q \geq 1$, $p+q = n$. The Szegő map S is a unitary operator from $W_{p,q-1}^{1,2}$ onto the space $\mathcal{H}^{p,q}$ of L^2 -harmonic (p, q) -forms.

Proof. Using [8], Theorem 4.9 and Lemma 4.2 we can compute easily the norm of $S(\gamma)$ in the $L^2(B, \wedge^{p,q})$ -space of (p, q) -forms on the unit ball,

$$(S(\gamma), S(\gamma))_{L^2(B, \wedge^{p,q})} = \frac{m!l!}{(n)_{m+l+2}} b^2(a+1)(b+1),$$

and by Lemma 3.2 above

$$(\gamma, \gamma)_{L^2(\mathcal{Q}^{p,q-1})} = \frac{m!l!}{(n)_{m+l+2}} b(b+1).$$

However

$$-(T + \bar{T})_* \gamma = ((m+q) - (l+p))\gamma = (b-a)\gamma$$

by direct computation, and

$$\partial_{\mathcal{Q}} \partial_{\mathcal{Q}} \gamma = a(b+1),$$

by Theorem 3.3. We have then

$$(\partial_{\mathcal{Q}} \partial_{\mathcal{Q}} - (T + \bar{T})_*) \gamma, \gamma = (S(\gamma), S(\gamma)),$$

completing the proof. $\square$

Remark 4.3. Theorem 4.2 needs to be modified slightly for $(p, q) = (n, 0)$ or $(p, q) = (0, n)$. Consider the case of $(p, q) = (n, 0)$. The space of harmonic $(n, 0)$ -forms are the Bergman space of holomorphic function $f(z)$ square integrable with respect to the Lebesgue measure with reproducing kernel $(1 - \langle z, w \rangle)^{-(n+1)}$, viewed as $(n, 0)$ -forms $f(z)(dz)^n$ where $(dz)^n$ stands for $dz_1 \wedge \cdots \wedge dz_n$. Consider the space of C^∞ -functions on S with the action of G :

$$\pi(g) : f(w) \mapsto f(g^{-1}w) J_{g^{-1}}(w)^{\frac{n}{n+1}}.$$

The Szegő map from functions on the sphere S to the Bergman space is then

$$(4.1) \quad Sf(z) = \int_S \frac{1}{(1 - \langle z, w \rangle)^{n+1}} f(w) d\sigma(w).$$

The L^2 -space on S is decomposed under $U(n)$ as a direct sum of irreducible subspaces with highest weights $(m, 0, \dots, 0, -l)$, see [10]. Let f be in the subspace with weight $(0, \dots, 0, -l)$, the Bergman space norm of Sf is given by

$$(4.2) \quad \|Sf\|^2_{(-\pi(T + \bar{T})f, f)_{L^2(S)}}$$

and thus extends to a unitary operator from the corresponding Sobolev space into the Bergman space. Here $T + \bar{T}$ is viewed as an element in the complexification of the Lie algebra of $U(n)$ with the induced action. This can be proved by a direct computation. The above scalar Szegő map is the same as the Szegő map in Theorems 4.1 and 4.2; indeed the area element is, writing

$$d\sigma(w) = \bar{\tau} \wedge (d\bar{\tau})^n = (d\bar{w})^n \wedge \left(\sum_{j=1}^n (-1)^{j-1} w_j dw_1 \wedge \cdots \widehat{dw_j} \cdots \wedge dw_n \right),$$

and identifying $Sf(z)$ in (4.1) as $Sf(z)(dz)^n$ we can write (4.1) as

$$Sf(z)(dz)^n = \int_S \frac{(dz)^n \wedge (d\bar{w})^n}{(1 - \langle z, w \rangle)^{(n+1)}} \wedge f(w) \left(\sum_{j=1}^n (-1)^{j-1} w_j dw_1 \wedge \cdots \widehat{dw_j} \cdots \wedge dw_n \right).$$

So if we identify the functions $f(w)$ on the boundary with $(n-1, 0)$ -form $F(w) = f(w)(\sum_{j=1}^n (-1)^{j-1} w_j dw_1 \wedge \cdots \widehat{dw_j} \cdots \wedge dw_n)$; the scalar Szegő map is then a map on forms

$$S(F)(z) \int_S \frac{(dz)^n \wedge (d\bar{w})^n}{(1 - \langle z, w \rangle)^{(n+1)}} \wedge F(w),$$

the integral kernel being then the (n, n) -form $(1 - \langle z, w \rangle)^{-(n+1)} (dz)^n \wedge (d\bar{w})^n$. So (4.2) takes the form

$$(4.3) \quad \|SF\|^2(-(T + \bar{T})_* F, F)_{L^2(S)}$$

with $(T + \bar{T})_*$ the induced action on differential forms of the vector field.

REFERENCES

- [1] T. Branson, *The functional determinant*, Tech. Report 4, Seoul National Univeristy, 1993.
- [2] ———, *Sharp inequalities, the functional determinant, and the complementary series*, Trans. Amer. Math. Soc. **347** (1995), 3671–3742.
- [3] Z. Chen, *Séries complémentaires des groupes de Lorentz et KK-théorie. (French) [Complementary series of Lorentz groups and KK-theory]*, J. Funct. Anal. **137** (1996), 76–96.
- [4] H. Donnelly and C. Fefferman, *L^2 -cohomology and index theorem for the Bergman metric*, Ann. Math **118** (1983), 593–618.
- [5] G. Folland, *The tangential Cauchy-Riemann complex on the sphere*, Trans. Amer. Math. Soc. **171** (1972), 83–133.
- [6] P.-Y. Gaillard, *Transformation de Poisson de Formes Differentielles. Le Cas de l'Espace Hyperbolique*, Comm. Math. Helv. **61** (1986), 581–616.
- [7] P. Griffiths and J. Harris, *Principles of algebraic geometry*, Wiley, New York, 1978.
- [8] P. Julg and G. Kasparov, *Operator K-theory for the group $SU(n, 1)$* , J. Reine Angew. Math. **463** (1995), 99–152.
- [9] J. Lott, *Invariant current on limit sets*, Comm. Math. Helv. **75** (2000), 319–350.
- [10] W. Rudin, *Function theory in the unit ball of $\mathbb{C}^n$* , Springer-Verlag, 1980.
- [11] D. P. Zelobenko, *Compact Lie groups and their representations*, Amer. Math. Soc., 1973.

BØ: DEPARTMENT OF MATHEMATICS, AARHUS UNIVERSITY, DENMARK

GZ: DEPARTMENT OF MATHEMATICS, CHALMERS UNIVERSITY OF TECHNOLOGY AND GOTHENBURG UNIVERSITY, GOTHENBURG, SWEDEN

E-mail address: orsted@imada.sdu.dk, genkai@math.chalmers.se